arXiv:1607.04768v1 [math.CO] 16 Jul 2016

HOFFMANN-OSTENHOF'S CONJECTURE FOR TRACEABLE CUBIC GRAPHS

F. Abdolhosseini[a], S. Akbari[b], H. Hashemi[a], M.S. Moradian[a]

[a] *Department of Computer Engineering, Sharif University of Technology, Tehran, Iran*
[b] *Department of Mathematical Sciences, Sharif University of Technology, Tehran, Iran*

Abstract. It was conjectured by Hoffmann-Ostenhof that the edge set of every connected cubic graph can be decomposed into a spanning tree, a matching and a family of cycles. In this paper, we show that this conjecture holds for traceable cubic graphs.

keywords: Cubic graph, Hoffmann-Ostenhof's Conjecture, Traceable
AMS Subject Classification: 05C45, 05C70

1. Introduction

Let G be a simple undirected graph with the *vertex set* $V(G)$ and the *edge set* $E(G)$. A vertex with degree one is called a *pendant vertex*. The distance between the vertices u and v in graph G is denoted by $d_G(u, v)$. A cycle C is called *chordless* if C has no *cycle chord* (that is an edge not in the edge set of C whose endpoints lie on the vertices of C). The *Induced subgraph* on vertex set S is denoted by $\langle S\rangle$. A path that starts in v and ends in u is denoted by $\widehat{vu}$. A *traceable* graph is a graph that possesses a Hamiltonian path. In a graph G, we say that a cycle C is *formed by the path* Q if $|E(C) \setminus E(Q)| = 1$. So every vertex of C belongs to $V(Q)$.

In 2011 the following conjecture was proposed:

Conjecture A. *(Hoffmann-Ostenhof [4]) Let G be a connected cubic graph. Then G has a decomposition into a spanning tree, a matching and a family of cycles.*

Conjecture A also appears in Problem 516 [3]. There are a few partial results known for Conjecture A. Kostochka [5] noticed that the Petersen graph, the prisms over cycles, and many other graphs have a decomposition desired in Conjecture A. Ozeki and Ye [6] proved that the conjecture holds for 3-connected cubic plane graphs. Furthermore, it was proved by Bachstein [2] that Conjecture A is true for every 3-connected cubic graph embedded in torus or Klein-bottle. Akbari, Jensen and Siggers [1, Theorem 9] showed that Conjecture A is true for Hamiltonian cubic graphs.

In this paper, we show that Conjecture A holds for traceable cubic graphs.

2. Results

Before proving the main result, we need the following lemma.

s_akbari@sharif.edu, {abdolhosseini, hohashemi, sadramoradian}@ce.sharif.edu.

Lemma 1. *Let G be a cubic graph. Suppose that $V(G)$ can be partitioned into a tree T and finitely many cycles such that there is no edge between any pair of cycles (not necessarily distinct cycles), and every pendant vertex of T is adjacent to at least one vertex of a cycle. Then, Conjecture A holds for G.*

Proof. By assumption, every vertex of each cycle in the partition is adjacent to exactly one vertex of T. Call the set of all edges with one endpoint in a cycle and another endpoint in T by Q. Clearly, the induced subgraph on $E(T) \cup Q$ is a spanning tree of G. We call it T'. Note that every edge between a pendant vertex of T and the union of cycles in the partition is also contained in T'. Thus, every pendant vertex of T' is contained in a cycle of the partition. Now, consider the graph $H = G \setminus E(T')$. For every $v \in V(T)$, $d_H(v) \leq 1$. So Conjecture A holds for G. □

Remark 1. Let C be a cycle formed by the path Q. Then clearly there exists a chordless cycle formed by Q.

Now, we are in a position to prove the main result.

Theorem 2. *Conjecture A holds for traceable cubic graphs.*

Proof. Let G be a traceable cubic graph and $P : v_1, \ldots, v_n$ be a Hamiltonian path in G. By [1, Theorem 9], Conjecture A holds for $v_1v_n \in E(G)$. Thus we can assume that $v_1v_n \notin E(G)$. Let $v_1v_j, v_1v_{j'}, v_iv_n, v_{i'}v_n \in E(G) \setminus E(P)$ and $j' < j < n$, $1 < i < i'$. Two cases can occur:

Case 1. Assume that $i < j$. Consider the following graph in Figure 1 in which the thick edges denote the path P. Call the three paths between v_j and v_i, from the left to the right, by P_1, P_2 and P_3, respectively (note that P_1 contains the edge e' and P_3 contains the edge e).



FIGURE 1. Paths P_1, P_2 and P_3

If P_2 has order 2, then G is Hamiltonian and so by [1, Theorem 9] Conjecture A holds. Thus we can assume that P_1, P_2 and P_3 have order at least 3. Now, consider the following subcases:

Subcase 1. There is no edge between $V(P_r)$ and $V(P_s)$ for $1 \leq r < s \leq 3$. Since every vertex of P_i has degree 3 for every i, by Remark 1 there are two chordless cycles C_1 and C_2 formed by P_1 and P_2, respectively. Define a tree T with the edge set

$$E\Big(\langle V(G) \setminus \big(V(C_1) \cup V(C_2)\big)\rangle\Big) \bigcap \big(\bigcup_{i=1}^{3} E(P_i)\big).$$

Now, apply Lemma 1 for the partition $\{T, C_1, C_2\}$.

Subcase 2. There exists at least one edge between some P_r and P_s, $r < s$. With no loss of generality, assume that $r = 1$ and $s = 2$. Suppose that $ab \in E(G)$, where $a \in V(P_1)$, $b \in V(P_2)$ and $d_{P_1}(v_j, a) + d_{P_2}(v_j, b)$ is minimum.



FIGURE 2. The edge ab between P_1 and P_2

Three cases occur:

(a) There is no chordless cycle formed by either of the paths $\overset{\frown}{v_j a}$ or $\overset{\frown}{v_j b}$. Let C be the chordless cycle $\overset{\frown}{v_j a} ab \overset{\frown}{v_j}$. Define T with the edge set

$$E\Big(\langle V(G) \setminus V(C)\rangle\Big) \bigcap \big(\bigcup_{i=1}^{3} E(P_i)\big).$$

Now, apply Lemma 1 for the partition $\{T, C\}$.

(b) There are two chordless cycles, say C_1 and C_2, respectively formed by the paths $\overset{\frown}{v_j a}$ and $\overset{\frown}{v_j b}$. Now, consider the partition C_1, C_2 and the tree induced on the following edges,

$$E\Big(\langle V(G) \setminus \big(V(C_1) \cup V(C_2)\big)\rangle\Big) \bigcap E\Big(\bigcup_{i=1}^{3} P_i\Big),$$

and apply Lemma 1.

(c) With no loss of generality, there exists a chordless cycle formed by the path $\overset{\frown}{v_j a}$ and there is no chordless cycle formed by the path $\overset{\frown}{v_j b}$. First, suppose that for every chordless cycle C_t on $\overset{\frown}{v_j a}$, at least one of the vertices of C_t is adjacent to a vertex in $V(G) \setminus V(P_1)$. We call one of the edges with

one end in C_t and other endpoint in $V(G)\setminus V(P_1)$ by e_t. Let $v_j = w_0, w_1, \ldots, w_l = a$ be all vertices of the path $\overset{\frown}{v_j a}$ in P_1. Choose the shortest path $w_0 w_{i_1} w_{i_2} \ldots w_l$ such that $0 < i_1 < i_2 < \cdots < l$. Define a tree T whose edge set is the thin edges in Figure 3.

Call the cycle $w_0 w_{i_1} \ldots w_l \overset{\frown}{bw_0}$ by C'. Now, by removing C', q vertex disjoint paths $Q_1, \ldots, Q_q$ which are contained in $\overset{\frown}{v_j a}$ remain. Note that there exists a path of order 2 in C' which by adding this path to Q_i we find a cycle C_{t_i}, for some i. Hence there exists an edge e_{t_i} connecting Q_i to $V(G)\setminus V(P_1)$. Now, we define a tree T whose the edge set is,

$$\Big(E\Big(\langle V(G)\setminus V(C')\rangle\Big)\bigcap\Big(\bigcup_{i=1}^{3}E(P_i)\Big)\Big)\bigcup\Big(\{e_{t_i}\mid 1\le i\le q\}\Big).$$

Apply Lemma 1 for the partition $\{T, C'\}$.



FIGURE 3. The cycle C' and the tree T

Next, assume that there exists a cycle C_1 formed by $\overset{\frown}{v_j a}$ such that none of the vertices of C_1 is adjacent to $V(G)\setminus V(P_1)$. Choose the smallest cycle with this property. Obviously, this cycle is chordless. Now, three cases can be considered:

(i) There exists a cycle C_2 formed by P_2 or P_3. Define the partition C_1, C_2 and a tree with the following edge set,

$$E\Big(\langle V(G)\setminus\big(V(C_1)\cup V(C_2)\big)\rangle\Big)\bigcap\Big(\bigcup_{i=1}^{3}E(P_i)\Big),$$

and apply Lemma 1.

(ii) There is no chordless cycle formed by P_2 and by P_3, and there is at least one edge between $V(P_2)$ and $V(P_3)$. Let $ab\in E(G)$, $a\in V(P_2)$ and $b\in V(P_3)$ and moreover $d_{P_2}(v_j,a)+d_{P_3}(v_j,b)$ is minimum. Notice that the cycle $\overset{\frown}{v_j ab v_j}$ is chordless. Let us call this cycle by C_2. Now, define the partition C_2 and a tree with the following edge set,

$$E\Big(\langle V(G)\setminus V(C_2)\rangle\Big)\bigcap\Big(\bigcup_{i=1}^{3}E(P_i)\Big),$$

and apply Lemma 1.

(iii) There is no chordless cycle formed by P_2 and by P_3, and there is no edge between $V(P_2)$ and $V(P_3)$. Let C_2 be the cycle consisting of two paths P_2 and P_3. Define the partition C_2 and a tree with the following edge set,

$$E\Big(\langle V(G) \setminus V(C_2)\rangle\Big) \bigcap \Big(\bigcup_{i=1}^{3} E(P_i)\Big),$$

and apply Lemma 1.

Case 2. Assume that $j < i$ for all Hamiltonian paths. Among all Hamiltonian paths consider the path such that $i' - j'$ is maximum. Now, three cases can be considered:

Subcase 1. There is no $s < j'$ and $t > i'$ such that $v_s v_t \in E(G)$. By Remark 1 there are two chordless cycles C_1 and C_2, respectively formed by the paths $v_1 v_{j'}$ and $v_{i'} v_n$. By assumption there is no edge xy, where $x \in V(C_1)$ and $y \in V(C_2)$. Define a tree T with the edge set:

$$E\Big(\langle V(G) \setminus \big(V(C_1) \cup V(C_2)\big)\rangle\Big) \bigcap \Big(E(P) \cup \{v_{i'} v_n, v_{j'} v_1\}\Big).$$

Now, apply Lemma 1 for the partition $\{T, C_1, C_2\}$.

Subcase 2. There are at least four indices $s, s' < j$ and $t, t' > i$ such that $v_s v_t, v_{s'} v_{t'} \in E(G)$. Choose four indices $g, h < j$ and $e, f > i$ such that $v_h v_e, v_g v_f \in E(G)$ and $|g - h| + |e - f|$ is minimum.



FIGURE 4. Two edges $v_h v_e$ and $v_g v_f$

Three cases can be considered:

(a) There is no chordless cycle formed by $v_g\widehat{v_h}$ and by $v_e\widehat{v_f}$.
Consider the cycle $v_g\widehat{v_h}v_e\widehat{v_f}\ v_g$ and call it C. Now, define a tree T with the edge set,

$$E\Big(\langle V(G) \setminus V(C)\rangle\Big) \bigcap \Big(E(P) \cup \{v_1 v_j, v_i v_n\}\Big),$$

apply Lemma 1 for the partition $\{T, C\}$.

(b) With no loss of generality, there exists a chordless cycle formed by $v_e\widehat{v_f}$ and there is no chordless cycle formed by the path $v_g\widehat{v_h}$. First suppose that there is a chordless cycle C_1 formed by $v_e\widehat{v_f}$ such that there is no edge between $V(C_1)$ and $\{v_1, \ldots, v_j\}$. By Remark 1, there exists a chordless cycle C_2 formed by $v_1\widehat{v_j}$. By assumption there is no edge between $V(C_1)$ and $V(C_2)$. Now, define a tree T with the edge set,

$$E\Big(\langle V(G)\setminus\big(V(C_1)\cup V(C_2)\big)\rangle\Big)\bigcap\Big(E(P)\cup\{v_1v_j,v_iv_n\}\Big),$$

and apply Lemma 1 for the partition $\{T,C_1,C_2\}$.

Next assume that for every cycle C_r formed by $v_e\widehat{v}_f$, there are two vertices $x_r\in V(C_r)$ and $y_r\in\{v_1,\dots,v_j\}$ such that $x_ry_r\in E(G)$. Let $v_e=w_0,w_1,\dots,w_l=v_f$ be all vertices of the path $v_e\widehat{v}_f$ in P. Choose the shortest path $w_0w_{i_1}w_{i_2}\dots w_l$ such that $0<i_1<i_2<\cdots<l$. Consider the cycle $w_0w_{i_1}\dots w_l\ v_g\widehat{v}_h$ and call it C. Now, by removing C, q vertex disjoint paths $Q_1,\dots,Q_q$ which are contained in $v_e\widehat{v}_f$ remain. Note that there exists a path of order 2 in C which by adding this path to Q_i we find a cycle C_{r_i}, for some i. Hence there exists an edge $x_{r_i}y_{r_i}$ connecting Q_i to $V(G)\setminus V(v_e\widehat{v}_f)$. We define a tree T whose edge set is the edges,

$$E\Big(\langle V(G)\setminus V(C)\rangle\Big)\bigcap\Big(E(P)\cup\{v_1v_j,v_iv_n\}\cup\big\{x_{r_i}y_{r_i}\mid 1\le i\le q\big\}\Big),$$

then apply Lemma 1 on the partition $\{T,C\}$.



FIGURE 5. The tree T and the shortest path $w_0w_{i_1}\dots w_l$

(c) There are at least two chordless cycles, say C_1 and C_2 formed by the paths $v_g\widehat{v}_h$ and $v_e\widehat{v}_f$, respectively. Since $|g-h|+|e-f|$ is minimum, there is no edge $xy\in E(G)$ with $x\in V(C_1)$ and $y\in V(C_2)$. Now, define a tree T with the edge set,

$$E\Big(\langle V(G)\setminus\big(V(C_1)\cup V(C_2)\big)\rangle\Big)\bigcap\Big(E(P)\cup\{v_1v_j,v_iv_n\}\Big),$$

and apply Lemma 1 for the partition $\{T,C_1,C_2\}$.

Subcase 3. There exist exactly two indices s,t, $s<j'<i'<t$ such that $v_sv_t\in E(G)$ and there are no two other indices s',t' such that $s'<j<i<t'$ and $v_{s'}v_{t'}\in E(G)$. We can assume that there is no cycle formed by $v_{s+1}\widehat{v}_j$ or $v_i\widehat{v}_{t-1}$, to see this by symmetry consider a cycle C formed by $v_{s+1}\widehat{v}_j$. By Remark 1 there exist chordless cycles C_1 formed by $v_{s+1}\widehat{v}_j$ and C_2 formed by $v_i\widehat{v}_n$. By assumption v_sv_t is the only edge such that $s<j$ and $t>i$. Therefore, there is no edge between $V(C_1)$ and $V(C_2)$. Now, let T be a tree defined by the edge set,

$$E\Big(\langle V(G)\setminus\big(V(C_1)\cup V(C_2)\big)\rangle\Big)\bigcap\Big(E(P)\cup\{v_1v_j,v_iv_n\}\Big),$$

and apply Lemma 1 for the partition $\{T,\,C_1,\,C_2\}$.

Furthermore, we can also assume that either $s\neq j'-1$ or $t\neq i'+1$, otherwise we have the Hamiltonian cycle $v_1\widehat{v}_sv_t\widehat{v}_nv_{i'}\widehat{v}_{j'}\,v_1$ and by [1, Theorem 9] Conjecture A holds.

By symmetry, suppose that $s\neq j'-1$. Let v_k be the vertex adjacent to $v_{j'-1}$, and $k\notin\{j'-2,j'\}$. It can be shown that $k>j'-1$, since otherwise by considering the Hamiltonian

path $P' : v_{k+1}\widehat{v}_{j'-1}v_k\widehat{v}_1v_{j'}\widehat{v}_n$, the new $i' - j'$ is greater than the old one and this contradicts our assumption about P in the Case 2.

We know that $j' < k < i$. Moreover, the fact that $v_{s+1}\widehat{v}_j$ does not form a cycle contradicts the case that $j' < k \leq j$. So $j < k < i$. Consider two cycles C_1 and C_2, respectively with the vertices $v_1\ v_{j'}\widehat{v}_j\ v_1$ and $v_n\ v_{i'}\widehat{v}_i\ v_n$. The cycles C_1 and C_2 are chordless, otherwise there exist cycles formed by the paths $v_{s+1}\widehat{v}_j$ or $v_i\widehat{v}_{t-1}$. Now, define a tree T with the edge set

$$E\Big(\langle V(G) \setminus \big(V(C_1) \cup V(C_2)\big)\rangle\Big) \bigcap \Big(E(P) \cup \{v_sv_t, v_kv_{j'-1}\}\Big),$$

and apply Lemma 1 for the partition $\{T, C_1, C_2\}$.

□

Remark 2. Indeed, in the proof of the previous theorem we showed a stronger result, that is, for every traceable cubic graph there is a decomposition with at most two cycles.

References

[1] S. Akbari, T.R. Jensen, M. Siggers, Decomposition of graphs into trees, forests, and regular subgraphs, Discrete Math. 338 (2015) no.8, 1322-1327.

[2] A.C. Bachstein, Decomposition of Cubic Graphs on the Torus and Klein Bottle, A Thesis Presented to the Faculty of the Department of Mathematical Sciences Middle Tennessee State University, 2015.

[3] P.J. Cameron, Research problems from the BCC22, Discrete Math. 311 (2011) 1074-1083.

[4] A. Hoffmann-Ostenhof, Nowhere-zero flows and structures in cubic graphs, Ph.D. dissertation, University at Wien, 2011.

[5] A. Kostochka, Spanning trees in 3-regular graphs, REGS in Combinatorics, University of Illinois at Urbana-Champaign, (2009). http://www.math.uiuc.edu/~west/regs/span3reg.html

[6] K. Ozeki, D. Ye, Decomposing plane cubic graphs, European J. Combin. 52 (2016), Part A, 40-46.